STM Knowledge Organiser Year: 8 Subject: Maths

plotted.

Core Know	ledge	1		
Topic/Skill	Definition/Tips	Example		
1. Coordinates	Written in pairs. The first term is the x-coordinate (movement across). The second term is the y-coordinate (movement up or down)	A: (4,7) B: (-6,-3)		
2. Midpoint of	Method 1: add the x coordinates and	Find the midpoint between (2,1) and		
a Line	divide by 2, add the y coordinates and divide by 2	(6,9)		
		$\frac{2+6}{2} = 4$ and $\frac{1+9}{2} = 5$		
	Method 2: Sketch the line and find the	2 2		
	values half way between the two x and two y values.	So, the midpoint is (4,5)		
3. Linear	Straight line graph.	Example:		
Graph		Other		
	The general equation of a linear graph is	examples:		
	y = mx + c	x = y		
	where m is the gradient and c is the y-	y = 4 $x = -2$		
	intercept.	y = 2x - 7		
		y + x = 10		
	The equation of a linear graph can contain an x-term , a y-term and a number .	2y - 4x = 12		
4. Plotting	Method 1: Table of Values	x -3 -2 -1 0 1 2 3		
Linear Graphs	Construct a table of values to calculate	y= x +3 0 1 2 3 4 5 6		
	coordinates.	y= x +3 0 1 2 3 4 5 6		
	Method 2: Gradient-Intercept Method	4		
	(use when the equation is in the form $y =$	33		
	mx + c)	$y = \frac{3}{2}x + 1$ 3		
	1. Plots the y-intercept	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \		
	2. Using the gradient, plot a second point.	0 2		
	3. Draw a line through the two points	3 -2 -1 0 1 2 3		
	plotted.			
	Method 3: Cover-Up Method (use when	97		
	the equation is in the form $ax + by = c$)	8 7		
	1. Cover the x term and solve the resulting	5		
	equation. Plot this on the $x - axis$.	3		
	2. Cover the y term and solve the resulting	1		
	equation. Plot this on the $y - axis$. 3. Draw a line through the two points	$3 \cdot 2 \cdot 10^{10} \cdot 12 \cdot 3 \cdot 10^{10} \cdot 12 \cdot 10^{10} \cdot 10$		
	2. Draw a fine unough the two points			

STM Knowledge Organiser Year: 8 Subject: Maths

Core Knowledge

Core Know	icusc	
5. Gradient	The gradient of a line is how steep it is.	Gradient = 4/2 = 2
	Gradient = $\frac{Change \ in \ y}{Change \ in \ x} = \frac{Rise}{Run}$	Gradient = -3/1 =-3
	The gradient can be positive (sloping upwards) or negative (sloping downwards)	1
6. Finding the Equation of a Line given a point and a gradient	Substitute in the gradient (m) and point (x,y) in to the equation $y = mx + c$ and solve for c.	Find the equation of the line with gradient 4 passing through (2,7). $y = mx + c$ $7 = 4 \times 2 + c$ $c = -1$
7. Finding the Equation of a Line given two points	Use the two points to calculate the gradient. Then repeat the method above using the gradient and either of the points.	$y = 4x - 1$ Find the equation of the line passing through (6,11) and (2,3) $m = \frac{11 - 3}{6 - 2} = 2$ $y = mx + c$ $11 = 2 \times 6 + c$ $c = -1$ $y = 2x - 1$
8. Parallel Lines	If two lines are parallel , they will have the same gradient . The value of m will be the same for both lines.	$y = 2x - 1$ Are the lines $y = 3x - 1$ and $2y - 6x + 10 = 0$ parallel? Answer: Rearrange the second equation in to the form $y = mx + c$ $2y - 6x + 10 = 0 \rightarrow y = 3x - 5$ Since the two gradients are equal (3), the lines are parallel.
9. Perpendicular Lines	If two lines are perpendicular , the product of their gradients will always equal -1 . The gradient of one line will be the negative reciprocal of the gradient of the other line. You may need to rearrange equations of lines to compare gradients (they need to be in the form $y = mx + c$)	Find the equation of the line perpendicular to $y = 3x + 2$ which passes through (6,5) Answer: As they are perpendicular, the gradient of the new line will be $-\frac{1}{3}$ as this is the negative reciprocal of 3. $y = mx + c$

STM Knowledge Organiser Year: 8 Subject: Maths Core Knowledge								
				$y = -\frac{1}{3}x + 7$				
			Or	3x + x - 7 = 0				
Links to sec	uences, proportion,	substitution, rate	of change,					